

## OXFORD PARTIAL (UNI) KNEE ARTHROPLASTY (UKA) DR. ORYHON

**PROCEDURE:** Partial Knee Arthroplasty      ☐ Right    ☐ Left    ☐ Bilateral

### GENERAL INSTRUCTIONS:

Use pain medication and/or ice packs when your knee hurts. It is advised (but not necessary) to take pain medication before you do your exercises and apply ice packs to your operative knee after exercises. Antithrombotic stockings (TED hose or compression stockings) are suggested for 4 weeks.

### KNEE PRECAUTIONS:

Do **not** use heat or heat creams on your operative knee/leg.

Do **not** lie with a pillow under your operative knee. You **may** place a pillow under the heel/ankle to promote straightening of your knee.

Do **not** twist, pivot or jump on operative knee/leg.

Do **not** immerse the operative knee under water (no baths/pools x6 weeks).

Do **not** drive until cleared to do so by Dr. Oryhon (typically 3-6 weeks).

### PHYSICAL THERAPY/REHAB:

Weight Bearing as Tolerated (WBAT) – use walker/cane as needed

Your inpatient rehab or home therapists will work on a UKA protocol, exercises include: ankle pumps, quad sets, straight leg raises, supine range of motion, seated range of motion, prone range of motion, terminal knee extension and gait training. *You should actively participate in your rehab and diligently do your prescribed exercises.*

*Note: Recovery from this procedure is often very rapid and in some cases can lead to “over-doing” activity in the early weeks which can result in increased inflammation and pain several weeks post-op after a period of feeling great; be patient and know that the body usually needs at least 6 weeks to heal after surgery.*

### WOUND CARE:

Unless otherwise informed, your incision is closed with absorbable suture – no suture or staple removal is necessary. Skin Glue (Dermabond) covers your incision to protect it while the skin heals – this will fall off with time.

Do **not** use ointments or creams on your operative knee.

Dry dressing change to incision if there is drainage (usually not needed past one week from surgery).

Shower is allowed (without covering wound) once wound is completely dry (no drainage on dressing).

### MEDICATIONS AND PRESCRIPTIONS:

Tylenol 1000mg every 8 hours for two weeks.

Additional pain medicine as prescribed. See “Medication and Education” section of your surgical packet.

Aspirin 325mg twice per day to thin the blood and prevent blood clots in **most** cases for 30 days.

Coumadin (Warfarin) is **sometimes** needed to thin the blood – the dose will be adjusted based on your blood tests.

An alternative oral medication (e.g. Xarelto, Eliquis) is sometimes used instead of Coumadin – no blood monitoring.

Colace as needed for constipation while on pain medicine.

Ferrous Sulfate 325mg each day (iron supplement to combat anemia) for 30 days.

**Also refer to the discharge medication reconciliation regarding your other medicines.**

### CLINIC FOLLOW-UP:

Dr. Oryhon will see you for follow-up 5-6 weeks after surgery

If you or your therapist have any concerns before the follow-up please call the office 847-381-0388.

Refer to our website [www.lakecookortho.com](http://www.lakecookortho.com) for full UKA knee rehab protocol and other information.